

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

costless water, the actual selling-price is $(\frac{10}{9})$ of $\frac{125}{60}$ of $100\% = 138\frac{8}{6}\%$. the actual rate per cent. of profit is $38\frac{8}{3}\%$.

- II. Solution by FRANK HORN, Columbia, Missouri, and Professor H. J. GAERTNER, Wilmington College, Wilmington, Ohio.
 - 1. 100% =apparent value.
 - 2. 125% = selling price of apparent value.
 - 3. 90% = 100% 10% = value of quantity sold for 125%.
- 4. .. $\frac{1_{80}^{2.5}\%}{1_{80}^{5.5}\%}$ = what 1% sells for. 5. $138_{9}^{8}\% = 100 \times \frac{1_{20}^{2.5}\%}{1_{20}^{5.5}\%}$ = real selling price. 6. 100% = true value.
 - - 7. $38\frac{8}{9}\% = 138\frac{8}{9}\% 100\% = \text{rate of gain.}$
 - ... The actual rate of gain is 38%.
- 34. A chain 100m long, weighing 14 oz. to the foot, is suspended from points on a level 80m apart. What is the sag, the batter at the ends, and the horizontal tension? [From Wentworth & Hill's High School Arithmetic.]

Solution by B. F. FINKEL, A. M., Professor of Mathematics and Physics in Kidder Institute. Kidder Missouri

The form of the chain fulfilling the conditions of the problem is the curve known as the catenary. Let B and C be the points of suspension of the chain, E any point in the chain, AL=x, EL=y.

Let AE=s and w=45.93oz, the weight of a metre of length of the chain.

Then ws=the weight of the portion AE=the load suspended at E, or the vertical tension at E. Let aw = the horizon-

tal tension at A, the weight of a units of length. Let EF be a tangent at E; then if EF represents the tension at E, EI and IF will represent the horizontal and vertical tensions respectively, at E.

Hence,
$$\frac{dy}{dx} = \frac{FI}{EI} = \frac{ws}{wa} = \frac{s}{a} \dots (1)$$
. But $ds = \sqrt{(dy^2 + dx^2)}$. $\therefore dy = \sqrt{(ds^2 - dx^2)}$. $s \neq a = \sqrt{(ds^2 - dx^2)} \neq dx$, whence

$$\frac{dx}{ds} = \frac{a}{\sqrt{(a^2 + s^2)}}. \quad \therefore \quad x = a \int \frac{ds}{\sqrt{(a^2 + s^2)}}$$

 $=a \log_e (s+1/\overline{a^2+s^2})+c.$ Since x=0, when s=0, $c=-a \log a$.

$$\therefore x = a \log_e \left[(s + \sqrt{a^2 + s^2}) / a \right] \dots (2). \quad \text{From (2), we have}$$

$$s = \frac{a}{2} \left(e^{\frac{x}{a}} - e^{-\frac{x}{a}} \right) \dots (3). \quad \text{From (1) and (3)} \frac{dy}{dx} = \frac{1}{2} \left(e^{\frac{x}{a}} - e^{-\frac{x}{a}} \right).$$

$$\therefore y = \frac{a}{2} \left(e^{\frac{x}{a}} - e^{-\frac{x}{a}} \right) + c. \text{ Since } y = 0 \text{ when } x = 0, c = -a.$$

$$\therefore y = \frac{a}{2} \left(e^{\frac{x}{a}} - e^{-\frac{x}{a}} \right) - a \dots (4). \quad \text{From (3) and (4) we get } a = (s^2 - y^2) \ge 2y.$$

From (2), we easily get $x = a \log_{e}[(y+1/y^{2} - a^{2}) / a] = a \log_{e}[(s+1/s^{2} + a^{2})]$ $|x| = (s^2 - y^2) / 2y \log [(s+y) / (s-y)] \div \log_{10} e.$

 $\log x = \log(s+y) + \log(s-y) + \log[\log(s+y) - \log(s-y)] - \log y - \log[2\log_{10} e].$ $= \log(s+y) + \log(s-y) + \log[\log(s+y) - \log(s-y)] + \cos(y+0.0612).$

From this equation, since x=40m and s=50m, we find, by the Method of Double Position, the value of y=26.53m which is called the sag.

The tension at
$$A = wa = w\left(\frac{s^2 - y^2}{2y}\right) = 1559.78$$
 oz., and $\frac{FI}{EI} = \frac{a}{s}$

=.6797 the butter.

From the above equations we may obtain the four propositions as given in Wentworth and Hill's High School Arithmetic.

35. Froposed by B. F. FINKEL. Professor of Mathematics in Kidder Institute, Kidder, Missouri.

Between Sing-Sing and Tarry-Town. I met my worthy friend, John Brown, And seven daughters, riding nags, and every one had seven bags: In every bag were thirty cats, and every cat had forty rats, Besides a brood of fifty kittens. All but the nags were wearing mittens! Mittens, kittens-cats, rats-bags, nags-Browns, How many were met between the towns?

[From Muttoon's Common Arithmetic.]

Solution by FRANK HORN, Columbia, Missouri.

- 1. 8=number of Browns met.
- 2. $8=8\times1=$ number of nags.
- 3. $56=8\times7=$ number of bags.
- 4. $1680=30\times56=$ number of cats.
- II. $\frac{1}{3}$ 5. 67200=1680 × 40=number of rats.
 - 6. $84000 = 1680 \times 50 = \text{number of kittens}$.

 - 7. 167888=Browns+cats+rats+kittens
 8. 335776=167888×2=number of mittens worn provided that each person, cat, rat, and kitten wore one pair.
 - 9. 636616=Browns+nags+bags+cats+rats+mittens+kittens.
- III. ... The number of objects and persons met amounted to 636616.

Nore.—The result given in Mattoon's Arithmetic is 2184192. What interpretation aid Mr. Mattoon give to the problem?—Editor.

PROBLEMS

42. Proposed by F. P. MATZ, M. So. Ph. D., Professor of Mathematics and Astronomy in New Windsor College, New Windsor, Maryland.

If m=2ct, be the interest on M=109ct, for p=19 days, find the yearly rate per cent:

43. Proposed by B. F. BURLESON, Oneida Castle, New York.

A, in a scuffle, seized on $\frac{2}{3}$ of a parcel of sugar plums; B caught $\frac{3}{3}$ of it out of his hands, and U laid hold on $\frac{3}{10}$ more; D ran off with all A had left, except 1 which E afterwards secured slyly for himself; then A and C jointly